# Monkey IL-17A ELISA development kit

Product Code: 3520M-1A-6

## CONTENTS, development kit for 6 plates:

### Vial 1 (yellow top)

Monoclonal antibody MT241 (150 µl)

Concentration: 0.5 mg/ml

## Vial 2 (red top)

Biotinylated monoclonal antibody MT504 (80 µl)

Concentration: 0.5 mg/ml

## Vial 3 (white top)

Streptavidin-Alkaline Phosphatase (80 µl)

#### Vial 4

Recombinant human IL-17A standard (1 μg)

To ensure total recovery of stated quantity, vials have been overfilled.

#### **STORAGE:**

Shipped at ambient temperature. On arrival box 1 should be stored refrigerated at 4-8°C and box 2 should be stored frozen at -20°C.

## General

**Intended use:** For quantitative determination of monkey IL-17A in solution, e.g. cell culture supernatant.

**Serum/plasma samples:** Please note that cytokine determinations in serum/plasma require the use of ELISA diluent (product code: 3652-D2) for dilution of samples, standard and detection antibody. The diluent prevents false positive read-outs which may be caused by interference of heterophilic antibodies found in plasma and serum. Please contact Mabtech for further information.

**Specificity:** Native and recombinant human IL-17A and native IL-17A from rhesus and cynomolgus macaques. Inquire for reactivities with other monkey species.


**Reagents:** Antibodies are supplied in sterile-filtered (0.2  $\mu$ m) PBS with sodium azide (0.02%). Streptavidin-ALP is supplied in 0.1 M Tris buffer with 0.002% Kathon CG.

Standard range: 3-300 pg/ml

**Intra-assay variation:** < 4%

**Standard calibration:** 1 ng of supplied standard equals 12 U of 01/420 NIBSC\*-standard according to repeated calibrations. Calibration is batch-specific.

\*National Institute of Biological Standards and Control, UK.



# Guidelines for Monkey IL-17A ELISA

- **Day 1** 1. Coat a high protein binding ELISA plate with mAb MT241, diluted to 1 μg/ml in PBS, pH 7.4, by adding 100 μl/well. Incubate overnight at 4-8°C.
- **Day 2** 2. Wash twice with PBS (200 μl/well).
  - 3. Block plate by adding 200  $\mu$ l/well of PBS with 0.05% Tween 20 containing 0.1% BSA (incubation buffer). Incubate for 1 hour at room temperature.
  - 4. Wash five times with PBS containing 0.05% Tween.
  - 5. Prepare hIL-17A standard by reconstituting contents of vial 4 in 1 ml PBS with 1% BSA to a concentration of 1  $\mu$ g/ml. Leave at room temperature for 15 minutes and then vortex the tube. The stock solution should be used immediately or stored in aliquots at -20°C for future use. We recommend the aliquots not be refrozen after initial use. For the test, prepare dilutions of the stock using the standard range as a guideline.
  - 6. Add 100 μl/well of samples or standards diluted in incubation buffer or ELISA diluent for serum/plasma samples and incubate for 2 hours at room temperature.
  - 7. Wash as in step 4.
  - 8. Add 100 μl/well of mAb MT504-biotin at 0.5 μg/ml in incubation buffer or ELISA diluent for serum/plasma samples. Incubate for 1 hour at room temperature.
  - 9. Wash as in step 4.
  - 10. Add 100 μl/well of Streptavidin-ALP diluted 1:1000 in incubation buffer. Incubate for 1 hour at room temperature.
  - 11. Wash as in step 4.
  - 12. Add 100 μl/well of appropriate substrate solution e.g. p-nitrophenyl-phosphate (pNPP), available from Mabtech product code 3652-P10.
  - 13. Measure the optical density (405 nm for pNPP) in an ELISA reader after suitable developing time.



Developed and manufactured by MABTECH AB, Sweden, whose quality management system complies with the standards ISO 9001:2015 & ISO 13485:2016.





#### The products are for research use only.

MABTECH shall not be liable for the use or handling of the product or for consequential, special, indirect or incidental damages there from.

Mabtech AB (Head Office)

Sweden Tel: +46 8 716 27 00 mabtech@mabtech.com Mabtech, Inc.

USA Tel: +1 513 871-4500 mabtech.usa@mabtech.com